

TABEL 1: TECHNISCHE GEGEVENS OMTRENT HET HOOFDGEWAS OP DE PERCELEN TE BOTTELARE IN HET 1^E PROEFJAAR.

	Perceel 1	Perceel 2
Hoofdgewas + variëteit	Triticale (verschillende variëteiten)	Wintertarwe
Oogstdatum	02/08/2011	01/08/2011
Oogstmethode	met proefveldmaaidorser	met proefveldmaaidorser
Datum persen stro	07/08/2011	05/08/2011
Datum ophalen stro	07/08/2011	05/08/2011
Datum opmeten stoppelhoogte	09/08/2011	09/08/2011
Datum stoppelbewerking	10/08/2011	10/08/2011
Type stoppelbewerking	frezen	frezen

TABEL 2: TECHNISCHE GEGEVENS OMTRENT BODEM EN BEMESTING OP DE PERCELEN TE BOTTELARE IN HET 1^E PROEFJAAR.

	Perceel 1	Perceel 2
Datum grondbewerking	10/08/2011	10/08/2011
Beschrijving grondbewerking	frezen	frezen
Datum bemonstering standaard grondontleding	09/08/2011	09/08/2011
Datum mesttoediening	10/08/2011	10/08/2011
Methode mesttoediening	mestinjectie	mestinjectie
Type mest	varkensdrijfmest (zeugen)	varkensdrijfmest (zeugen)
Hoeveelheid toegediende mest (ton/ha) voor de bemestingstrappen 60N en 120N	Voor 60N: 14.5 ton/ha Voor 120N: 29 ton/ha	Voor 60N: 14.5 ton/ha Voor 120N: 29 ton/ha
Datum staalname toegediende mest	02/08/2011 (voor toediening, in mestput) 10/08/2011 (aan injecteur bij toedienen)	02/08/2011 (voor toediening, in mestput) 10/08/2011 (aan injecteur bij toedienen)

Beschrijving staalname toegediende mest	in mestput; mengstaal na mixen aan injecteur: mengstaal op verschillende tijdstippen (bij start injectie en bij uitloop)	in mestput; mengstaal na mixen aan injecteur: mengstaal op verschillende tijdstippen (bij start injectie en bij uitloop)
---	--	--

TABEL 3: TECHNISCHE GEGEVENS OMTRENT HET VANGGEWAS TE BOTTELARE IN HET 1^E PROEFJAAR (OP BEIDE PERCELEN).

Type vanggewas	Gele mosterd	Italiaans raaigras	Japanse haver	Gras/klaver
Datum 1e inzaai	10/08/2011	10/08/2011	10/08/2011	niet ingezaaid
Inzaaimethode (+ merknaam zaaimachine)	proefveldzaaimachine (breedte van 1,5 m)			niet ingezaaid
Zaadichtheid (per bestanddeel)	20 kg/ha	niet ingezaaid	75 kg/ha	niet ingezaaid
Naam variëteit(en)	Albatros	niet ingezaaid	Preta Comum	niet ingezaaid
Datum 2e inzaai	29/08/2011	niet ingezaaid	29/08/2011	niet ingezaaid
Inzaaimethode	proefveldzaaimachine (breedte van 1,5 m)			niet ingezaaid
Zaadichtheid (per bestanddeel)	20 kg/ha	niet ingezaaid	75 kg/ha	niet ingezaaid
Naam variëteit(en)	Albatros	niet ingezaaid	Preta Comum	niet ingezaaid
Oogstmethode	Haldrup (breedte van 1,5 m)			niet ingezaaid
Datum onkruidbestrijding braak	13/09/2011			niet ingezaaid

TABEL 4: TECHNISCHE GEGEVENS OMTRENT HET HOOFDGEWAS VOOR HET PERCEEL TE LEMBERGE IN HET 1^E PROEFJAAR.

Hoofdgewas + variëteit	Wintergerst
Oogstdatum	06/07/2011
Oogstmethode	pikdorsen
Datum persen stro	08/07/2011
Datum ophalen stro	08/07/2011
Datum opmeten stoppelhoogte	17/07/2011

Datum stoppelbewerking	17/07/2011
Type stoppelbewerking	klepelen

TABEL 5: TECHNISCHE GEGEVENS OMTRENT BODEM EN BEMESTING VOOR HET PERCEEL TE LEMBERGE IN HET 1^E PROEFJAAR.

Datum grondbewerking	03/08/2011
Beschrijving grondbewerking	Vastetandcultivator direct na mengmesttoepassing, rotoreg in 1 werkgang met zaaimachine
Datum bemonstering standaard grondontleding	02/08/2011
Datum mesttoediening	03/08/2011
Methode mesttoediening	Akkerbouwbemester Vervaet/ pijpen net bovengronds laten lopen en direct inwerken
Type mest	mestvarkensmengmest
Hoeveelheid toegediende mest (ton/ha) voor de bemestingstrappen 60N en 120N	Voor 60N: 7 ton Voor 120N: 14 ton
Datum staalname toegediende mest	03/08/2011
Beschrijving staalname toegediende mest	Opvangen bij stilstand machine tijdens toepassing

TABEL 6: TECHNISCHE GEGEVENS OMTRENT HET VANGGEWAS VOOR HET PERCEEL TE LEMBERGE IN HET 1^E PROEFJAAR.

Type vanggewas	Gele mosterd	Italiaans raaigras	Japanse haver	Gras/klaver
Datum 1e inzaai	04/08/2011	04/08/2011	04/08/2011	04/08/2011
Inzaaimethode (+ merknaam zaaimachine)	Graanzaaimachine 'Amazone' 3 m breed en rijafstand 10 cm			
Zaadichtheid (per bestanddeel)	25 kg/ha	45 kg/ha	75 kg/ha	42 kg/ha
Naam variëteit(en)	Salsa	Meltop	Preta Comum	Mengsel Engels raaigras ('Meltador' 15kg + 'Melpro' 15kg) + rode klaver ('Merviot' 8 kg) + witte klaver ('Merwi' 4 kg)

Datum 2e inzaai	31/08/2011	31/08/2011	31/08/2011	31/08/2011
Inzaaimethode	Graanzaaimachine 'Amazone' 3 m breed en rijafstand 10 cm			
Zaadichtheid (per bestanddeel)	25 kg/ha	45 kg/ha	75 kg/ha	42 kg/ha
Naam variëteit(en)	Salsa	Meltop	Preta comum	Mengsel Engels raaigras (Meltador 15kg + Melpro 15kg) + rode klaver ('Merviot' 8 kg) + witte klaver ('Merwi' 4 kg)
Oogstmethode	Haldrup (breedte van 1,4 m)			
Datum onkruidbestrijding braak	11/08/2011, herhaling op 14/10/2011			

TABEL 7: TECHNISCHE GEGEVENS OMTRENT HET HOOFDGEWAS VOOR HET PERCEEL TE RUKKELINGEN-LOON IN HET 1^E PROEFJAAR.

Hoofdgewas + variëteit	Wintertarwe
Oogstmethode	Maaidorser
Datum opmeten stoppelhoogte	17/08/11
Datum stoppelbewerking	17/08/11
Type stoppelbewerking	Vastetandcultivator

TABEL 8: TECHNISCHE GEGEVENS OMTRENT BODEM EN BEMESTING VOOR HET PERCEEL TE RUKKELINGEN-LOON IN HET 1^E PROEFJAAR.

Datum grondbewerking	17/08/11
Beschrijving grondbewerking	Na toedienen varkensdrijfmest bewerking met cultivator. Na zaai toerollen.
Datum bemonstering standaard grondontleding	29/09/11
Datum mesttoediening	17/08/11
Methode mesttoediening	Sleepslangen

Type mest	Varkensdrijfmest
Hoeveelheid toegediende mest (ton/ha) voor de bemestingstrappen 60N en 120N	Voor 60N: 8 ton/ha Voor 120N: 16 ton/ha
Datum staalname toegediende mest	17/08/11
Beschrijving staalname toegediende mest	Tweemaal staalname aan toestel

TABEL 9: TECHNISCHE GEGEVENS OMTRENT BODEM EN BEMESTING VOOR HET PERCEEL TE RUKKELINGEN-LOON IN HET 1^E PROEFJAAR.

Type vanggewas	Gele mosterd	Italiaans raaigras	Japane haver	Gras/klaver
Datum 1e inzaai	18/08/11	18/08/11	18/08/11	18/08/11
Zaadichtheid (per bestanddeel)	25 kg/ha	45 kg/ha	75 kg/ha	15kg/ha tetraploïd; 15 kg/ha diploid; 8 kg/ha rode klaver; 4 kg/ha witte klaver
Naam variëteit(en)	Salsa	Meltop (tetraploïd)		Meltador (tetraploïd); Melpro (diploid); Merviot: witte klaver
Datum 2e inzaai	31/08/11	31/08/11	31/08/11	31/08/11
Zaadichtheid (per bestanddeel)	25 kg/ha	45 kg/ha	75 kg/ha	15kg/ha tetraploïd; 15 kg/ha diploid; 8 kg/ha rode klaver; 4 kg/ha witte klaver
Naam variëteit(en)	Salsa	Meltop (tetraploïd)	-	Meltador (tetraploïd); Melpro (diploid); Merviot: witte klaver
Oogstmethode	Maaibalk, per veld opbrengst van 8 m ² bijeengeharkt Plukstaal voor analyse			Gras/klaver was onvoldoende ontwikkeld en kon niet worden geoogst.
Datum onkruidbestrijding braak	29/09/11			

TABEL 10: TECHNISCHE GEGEVENS OMTRENT HET HOOFDGEWAS VOOR HET PERCEEL TE SINT-DENIJS IN HET 1^E PROEFJAAR.

Hoofdgewas + variëteit	wintertarwe
Oogstdatum	03/08/11

Oogstmethode	dorsen
Datum persen stro	10/08/11
Datum ophalen stro	10/08/11
Datum opmeten stoppelhoogte	09/08/11 (50 metingen)

TABEL 11: TECHNISCHE GEGEVENS OMTRENT BODEM EN BEMESTING VOOR HET PERCEEL TE SINT-DENIJS IN HET 1^E PROEFJAAR.

Datum grondbewerking	12/08/11 (grondbewerking na mengmesttoediening; grondbewerking in de percelen "zaaitijdstip 1", "zaaitijdstip 2" en "braak"
Beschrijving grondbewerking	cultivator
Datum bemonstering standaard grondontleding	09/08/11
Datum mesttoediening	11/08/11
Methode mesttoediening	mestinjectie, 15 cm in de grond
Type mest	zeugendrijfmest
Hoeveelheid toegediende mest (ton/ha) voor de bemestingstrappen 60N en 120N	14 en 28
Datum staalname toegediende mest	11/08/11
Beschrijving staalname toegediende mest	aan tankwagen

TABEL 12: TECHNISCHE GEGEVENS OMTRENT HET VANGGEWAS VOOR HET PERCEEL TE SINT-DENIJS IN HET 1^E PROEFJAAR.

Type vanggewas	Gele mosterd	Italiaans raigras	Japane haver	Gras/klaver
Datum 1e inzaai	12/08/11	12/08/11	12/08/11	12/08/11
Inzaaimethode (+ merknaam zaaimachine)	Rotoreg Kuhn Zaaimachine Venta	Rotoreg Kuhn Zaaimachine Venta	Rotoreg Kuhn Zaaimachine Venta	Rotoreg Kuhn Zaaimachine Venta
Zaadichtheid (per bestanddeel)	25 kg/ha	45 kg/ha	75 kg/ha	42 kg/ha

Naam variëteit(en)	Salsa	Meltop	Mulata	Engels gras: Meltador en Melpro rode klaver: Merviot witte klaver: Merwi
Datum 2e inzaai	30/08/11	30/08/11	30/08/11	30/08/11
Inzaaimethode	Rotoreg Kuhn Zaaimachine Venta	Rotoreg Kuhn Zaaimachine Venta	Rotoreg Kuhn Zaaimachine Venta	Rotoreg Kuhn Zaaimachine Venta
Zaadichtheid (per bestanddeel)	25 kg/ha	45 kg/ha	75 kg/ha	42 kg/ha
Naam variëteit(en)	Salsa	Meltop	Mulata	Engels gras: Meltador en Melpro rode klaver: Merviot witte klaver: Merwi
Oogstmethode	cirkelmaaier	cirkelmaaier	Cirkelmaaier	- cirkelmaaier: zaaitijdstip 1 (behalve het veldje gras/klaver onbemest 1 ^e zaaitijdstip, parallel 3) - gazonmaaier: zaaitijdstip 2 en het veldje gras/klaver onbemest 1 ^e zaaitijdstip, parallel 3
Datum onkruidbestrijding braak	21/09/2011, herhaling op 27/10/2011			

TABEL 13: TEELTTECHNISCHE EN OPBRENGSTGEGEVENS VOOR HET VOORGEWAS EN TECHNISCHE GEGEVENS OMTRENT DE MESTTOEDIENING EN DE BODEMBEWERKING IN HET 2E PROEFJAAR.

	Bottelare	Merelbeke	Oostende	Ruk-Loon
<i>Voorgewas</i>	Triticale	Gerst (Zoom)	Tarwe (Lear)	Tarwe (Viscount)
<i>Zaaidatum</i>	24/10/11	1 ^e week 10/11	18/10/11	18/10/11
<i>N_{min} (0-90 cm) in voorjaar (kg N/ha)</i>	26	65,2	111,5	53,6
<i>Stikstofadvies (kg N werkzaam/ha)</i>	178	140	176	200
<i>Max. bemesting MAP4 wintergraan + nateelt (kg N werkzaam/ha)</i>	180	145	195	195
<i>Datum en hoeveelheid (kg N/ha) bemesting op</i>	F1: 23/03/12: 56 F2: 23/04/12: 56 F3: 09/05/12: 54	F1: 15/03/12: 54 F2: 24/04/12: 54	F1: 7/03/12: 68 F2: 14/04/12: 33 F3: 28/05/12: 49	27/03/12: 75 20/04/12: 40 19/05/12: 55

<i>voorgewas</i>	som: 166	som: 108	som:150	som: 170
<i>Type bemesting op voorgewas</i>	F1: vloeibare N F2: vloeibare N F3: NH ₄ NO ₃	F1: NH ₄ NO ₃ F2: NH ₄ NO ₃	F1: NH ₄ NO ₃ F2: vloeibare N F3: NH ₄ NO ₃	F1: vloeibare N F2: vloeibare N F3: NH ₄ NO ₃
<i>Oogstdatum</i>	12/08/12	24/07/12	12/08/12	14/08/12
<i>Graanopbrengst (ton/ha)</i>	10	6,6	9,2	9
<i>Stro-opbrengst (ton/ha)</i>	2,8	2,9	4,7	4
<i>Stoppellengte (cm)</i>	15	13,8	-	10 tot 15
<i>Datum mesttoediening</i>	21/08/12	31/07/12	21/08/12	18/08/12
<i>Methode mesttoediening</i>	Injectie	Sleepslangen	Sleepvoet injectie	Sleepslangen
<i>Datum bodembewerking</i>	22/08/2012	31/07/12	21/08/12	18/08/12
<i>Type bodembewerking</i>	Optrekken en rotoeggen	Vastetandcultivator	Vastetandcultivator	Tweemaal met cultivator

TABEL 14: VARIËTEITEN EN ZAAIDICHTHEDEN VOOR DE IN HET IN 2^e PROEFJAAR GEBRUIKTE VANGGEWASSEN (ALLE LOCATIES).

<i>Vanggewas</i>	Gele mosterd	Italiaans raaigras	Japanse haver	Grasklavermengsel
<i>Variëteit</i>	Salsa	Meltop	Pratex	Melpro/Meltador/ Merviot/Merwi
<i>Zaaidichtheid</i>	25 kg/ha	45 kg/ha	75 kg/ha	Engels raaigras: 30 kg/ha Rode klaver: 8 kg/ha Witte klaver: 4 kg/ha
<i>Rijafstand</i>	10,5 tot 15 cm	10,5 tot 15 cm	10,5 tot 15 cm	10,5 tot 15 cm

Bottelare, perceel 1, 1^e proefjaar


Behandelingen:

- 1: Braak
- 2: Gele mosterd 20 kg/ha
- 3: Italiaans raaigras 45 kg/ha
- 4: Japanse haver 75 kg/ha

N0: 0 kg N/ha

N60: 60 kg/ha

N120: 120 kg/ha



Bottelare, perceel 2, 1^e proefjaar


Behandelingen:

- 1: Braak
- 2: Gele mosterd 20 kg/ha
- 3: Italiaans raaigras 45 kg/ha
- 4: Japanse haver 75 kg/ha


N0: 0 kg N/ha

N60: 60 kg/ha

N120: 120 kg/ha



Lemberge, 1^e proefjaar




Sint-Denijs, 1^e proefjaar

Zandleembodem																																																																																																				
Voorvrucht: wintertarwe																																																																																																				
		1				2				3				4				5				1		5		4		2		3																																																																						
		braak				gele mosterd 25 kg/ha				it raai gras 45 kg/ha				japanse h. 75 kg/ha				gras klaver 42 kg/ha																																																																																		
		1 ^o zaai		2 ^o zaai		2 ^o zaai		1 ^o zaai		2 ^o zaai		2 ^o zaai		1 ^o zaai		2 ^o zaai		2 ^o zaai		1 ^o zaai		2 ^o zaai		2 ^o zaai		1 ^o zaai		2 ^o zaai																																																																								
		90 m										225 m										90 m																																																																														
		15 m										15 m										15 m																																																																														
		10 m																																																																																																		
		12 m																																																																																																		
		0										60										120																																																																														
36 m		60										120										0																																																																														
		120										60										120																																																																														
		0										60										120																																																																														
36 m		60										120										0																																																																														
		120										60										120																																																																														
		0										60										120																																																																														
		120										60										120																																																																														
		12 m																																																																																																		
		2 ^o zaai				1 ^o zaai				1 ^o zaai				2 ^o zaai				1 ^o zaai				2 ^o zaai				1 ^o zaai																																																																										
		4				5				1				2				3				4				1				2				3				5																																																														
		oogsten min 10 m ² /veldje																																																																																																		
0																																																																																																				
60																																																																																																				
120																																																																																																				
		101	104	119	122	107	110	125	128	113	202	229	214	211	226	220	205	208	223	60	203	230	215	212	227	221	206	209	224	120	201	228	213	210	225	219	204	207	222	0	326	311	314	329	302	320	305	308	323	120	412	427	403	421	406	409	424	430	415	120	325	310	313	328	301	319	304	307	322	60	411	426	402	420	405	408	423	429	414	60	327	312	315	330	303	321	306	309	324	0	410	425	401	419	404	407	422	428	413	0

Rukkelingen-Loon, 1^e proefjaar


De vier blokken vertegenwoordigen de vier herhalingen, de twee zaaitijdstippen zijn telkens aangegeven. De verschillende kleuren komen overeen met de behandeling zoals onderaan getoond in de legende; hoe donkerder de tint, hoe hoger het bemestingsniveau.



Legende

braak	mosterd	lt raai gras	Japanse Haver	gras-klover
-------	---------	--------------	---------------	-------------

Bottelare, 2e proefjaar



zaai 1 | zaai 2

zaai 1 | zaai 2



- 1: Braak
- 2: Gele mosterd 25 kg/ha
- 3: Italiaans raaigras 45 kg/ha
- 4: Japanse haver 75 kg/ha
- 5: Gras/klaver 42 kg/ha

Merelbeke, 2e proefjaar


15 m

- | |
|----------------------|
| 1 blanco |
| 2 gele mosterd |
| 3 italiaans raagrass |
| 4 Japanse haver |
| 5 gras / klaver |

par



Oostende, 2e proefjaar




0 EN
60 EN
120 EN

↑ zaairichting; per veldje 4 zaaimachines x 2,5 m breedte = 10 m breedte
→ richting drijfmesttoediening; per veldje 2 x 6 m (= breedte mestinjecteur) = 12 m breedte

Rukkelingen-Loon, 2e proefjaar

AARDAPPELEN

24 M



		braak		gele mosterd 20kg/ha		It raaigras 45kg/ha		Japanse Haver 75kg/ha		gras-klover 42 kg/ha	
		1e zaai	2e zaai	2e zaai	1e zaai	1e zaai	2e zaai	2e zaai	1e zaai	1e zaai	2e zaai
36m	0	6	12	18	24	30	36	42	48	54	
	60	5	11	17	23	29	35	41	47	53	
	120	4	10	16	22	28	34	40	46	52	
12m											
36m	60	3	9	15	21	27	33	39	45	51	
	0	2	8	14	20	26	32	38	44	50	
	120	1	7	13	19	25	31	37	43	49	
		2e zaai	1e zaai	1e zaai	2e zaai	2e zaai	1e zaai	1e zaai	2e zaai		
		Japanse Haver 75kg/ha		gras-klover		braak		mosterd 20k/ha		It raaigras 45kg/ha	

		braak		gras-klover 42 kg/ha		Japanse Haver 75kg/ha		gele mosterd 20kg/ha		It raaigras 45kg/ha	
		2e zaai	1e zaai	1e zaai	2e zaai	2e zaai	1e zaai	2e zaai	1e zaai	1e zaai	2e zaai
60	60	60	66	72	78	84	90	96	102	108	
	120	59	65	71	77	83	89	95	101	107	
	0	58	64	70	76	82	88	94	100	106	
120	57	63	69	75	81	87	93	99	105		
	60	56	62	68	74	80	86	92	98	104	
	0	55	61	67	73	79	85	91	97	103	
		1e zaai	2e zaai	2e zaai	1e zaai	1e zaai	2e zaai	2e zaai	1e zaai		
		Japanse Haver 75kg/ha		braak		mosterd 20k/ha		It raaigras 45kg/ha		gras-klover	

TABEL 1: STANDAARD GRONDONTLEDINGEN VAN DE PROEFVELDLOCATIES.

Locatie	proefjaar	textuur (manueel)	pH- KCl	OC (%)	P (mg/100g)	K (mg/100g)	Mg (mg/100g)	Ca (mg/100g)	Na (mg/100g)	Fe (mg/100g)	Mn (mg/100g)
Bottelare 1	2011-2012	lemig zand	5,0	1,0	28	13	3	51	<0,9	-	-
Bottelare 2	2011-2012	licht zandleem	5,7	0,8	23	15	13	61	<0,9	-	-
Lemberge	2011-2012	zandleem	5,6	1,4	25	16	15	85	<0,9	-	-
Rukkelingen-Loon	2011-2012	leem	6,8	1,2	19	22	17	257	3,8	-	-
Sint-Denijs	2011-2012	zandleem	6,8	1,07	28	19	14	258	<2,0	-	-
Bottelare	2012-2013	zand	5,2	1,23	27	10	5	52	<0,9	-	-
Merelbeke	2012-2013	zandleem	5,6	1,13	37	15	9	95	2,2	74	8,6
Oostende	2012-2013	klei	7,2	1,25	20	23	48	1126	4,3	-	-
Rukkelingen-Loon	2012-2013	leem	6,6	1,10	22	21	15	254	1,5	-	-

De bemonstering werd uitgevoerd in juli-augustus 2011 voor het 1e proefjaar en in juli-augustus 2012 voor het 2e proefjaar. Gehaltes aan fosfor, kalium, magnesium, calcium en natrium zijn weergegeven in mg/100g luchtdroge grond en werden gemeten in het A.L.-extract.

1. OVERZICHTSFOTO'S



FIGUUR 1: VELDPROEF TE BOTTELARE, 21 SEPTEMBER 2011. OP DE VOORGROND DE 2^E ZAAI VAN GELE MOSTERD (L) EN JAPANSE HAVER (R) OP BEMESTINGSTRAP 120N, DAARACHTER DEZELFDE VANGGEWASSEN OP DE 0N.



FIGUUR 2: VELDPROEF TE LEMBERGE, 14 SEPTEMBER 2011. OP DE VOORGROND DE 1^E ZAAI VAN ITALIAANS RAAIGRAS EN GELE MOSTERD OP BEMESTINGSTRAPPEN 120N – 60N – 0N VAN LINKS NAAR RECHTS.



FIGUUR 3: VELDPROEF TE RUKKELINGEN-LOON, 10 OKTOBER 2011. CENTRAAL IN BEELD DE MET TOTAALHERBICIDE BEHANDELDE BRAAKVELDJES.



FIGUUR 4: VELDPROEF TE SINT-DENIJS, 24 OKTOBER 2011. OP DE ACHTERGROND DE 1^E ZAAI VAN GELE MOSTERD OP BEMESTINGSTRAPPEN 0N – 120N – 60N VAN LINKS NAAR RECHTS.



FIGUUR 5: VELDPROEF TE BOTTELARE, 24 OKTOBER 2012. CENTRAAL IN BEELD DE 1^E ZAAI VAN JAPANSE HAVER OP BEMESTINGSTRAPPEN 0N – 60N – 120N VAN LINKS NAAR RECHTS.



FIGUUR 6: VELPPOEF TE MERELBEKE, 12 SEPTEMBER 2012. OP DE ACHTERGROND DE 1^E ZAAI VAN GELE MOSTERD OP BEMESTINGSTRAPPEN 120N – 60N – 0N VAN LINKS NAAR RECHTS.



FIGUUR 7: VELDPROEF TE MERELBEKE, 7 OKTOBER 2012. UITERST LINKS DE 1^E ZAAI VAN GELE MOSTERD, RECHTS DAARVAN DE 1^E ZAAI VAN JAPANSE HAVER. VERVOLGENS VAN LINKS NAAR RECHTS DE 2^E ZAAI VAN ITALIAANS RAAIGRAS, JAPANSE HAVER EN GELE MOSTERD.



FIGUUR 8: VELDPROEF TE OOSTENDE, 23 OKTOBER 2012. OP DE VOORGROND DE 1^E ZAAI VAN GELE MOSTERD OP DE 120N BEMESTINGSTRAP; DE VANGGEWASSEN ONTKIEMDEN LAAT EN ONTWIKKELDEN ZICH SLECHT DOOR DE DROOGTE BIJ EN NA INZAAI.



FIGUUR 9: VELDPROEF TE RUKKELINGEN-LOON, 18 OKTOBER 2012. OP DE VOORGROND DE 1^E ZAAI VAN JAPANSE HAVER.



FIGUUR 10: VELDPROEF TE RUKKELINGEN-LOON, 21 DECEMBER 2012. LINKS DE BRAAKPERCELEN, CENTRAAL DE GEKLEPELDE NIET-WINTERHARDE VANGGEWASSEN (GELE MOSTERD EN JAPANSE HAVER) EN ACHTERAAN DE GEKLEPELDE WINTERHARDE VANGGEWASSEN (ITALIAANS RAAIGRAS EN HET GRASKLAVERMENGSEL).



FIGUUR 11: BOTTELARE, 22 FEBRUARI 2013. VAN LINKS NAAR RECHTS: ITALIAANS RAAIGRAS, BRAAK, GELE MOSTERD, JAPANSE HAVER EN HET GRASKLAVERMENGSEL .

2. DETAILFOTO'S

In de volgende detailfoto's wordt eerst het onderscheid tussen vanggewassen op de 0N en de 60N bemestingstrap in beeld gebracht, nadien komt het verschil tussen beide zaaitijdstippen aan bod.



FIGUUR 12: LEMBERGE, 14 SEPTEMBER 2011. 1^E ZAAI VAN GELE MOSTERD OP 60N (L) EN 0N (R).



FIGUUR 13: LEMBERGE, 14 SEPTEMBER 2011. 1^E ZAAI VAN ITALIAANS RAAIGRAS OP 60N (L) EN 0N (R).



FIGUUR 14: LEMBERGE, 14 SEPTEMBER 2011. 1^E ZAAI VAN JAPANSE HAVER OP 60N (L) EN 0N (R).



FIGUUR 15: LEMBERGE, 14 SEPTEMBER 2011. 1^E ZAAI VAN HET GRASKLAVERMENGSEL OP 60N (L) EN 0N (R).



FIGUUR 16: RUKKELINGEN-LOON, 24 AUGUSTUS 2011. 1^E ZAAI VAN GELE MOSTERD 6 DAGEN NA INZAAI. DE VERSLEMPING DOOR TOEDOEN VAN DE STORM KORT NA INZAAI IS DUIDELIJK WAARNEEMBAAR.



FIGUUR 17: RUKKELINGEN-LOON, 10 OKTOBER 2011. 2^E ZAAI VAN GRASKLAVER (ENGELS RAAIGRAS + RODE KLAVER + WITTE KLAVER) OP 60N BEMESTINGSTRAP.



FIGUUR 18: SINT-DENIJS, 24 OKTOBER 2011. GELE MOSTERD 1^E ZAAI (R) EN 2^E ZAAI (L) OP DE 60N BEMESTINGSTRAP.



FIGUUR 19: RUKKELINGEN-LOON, 10 OKTOBER 2011. ITALIAANS RAAIGRAS 1^E ZAAI (L) EN 2^E ZAAI (R) OP DE 0N BEMESTINGSTRAP.




FIGUUR 20: BOTTELARE, 24 OKTOBER 2012. JAPANSE HAVER 1^E ZAAI (L) EN 2^E ZAAI (R) OP DE ON BEMESTINGSTRAP.




FIGUUR 21: SINT-DENIJS, 24 OKTOBER 2011. GRASKLAVER 1^E ZAAI (R) EN 2^E ZAAI (L) OP DE 60N BEMESTINGSTRAP.



BOVENGRONDSE STIKSTOFOPBRENGST BOTTELARE





BOVENGRONDSE STIKSTOFOPBRENGST LEMBERGE



BOVENGRONDSE STIKSTOFOPBRENGST RUKKELINGEN-LOON




De vanggewassen werden ondergewerkt op 15 januari 2012; er werd geen voorjaarsoogst meer uitgevoerd.




De vanggewassen werden ondergewerkt op 15 januari 2012; er werd geen voorjaarsoogst meer uitgevoerd.


BOVENGRONDSE STIKSTOFOPBRENGST SINT-DENIJS





BOVENGRONDSE STIKSTOFOPBRENGST BOTTELARE





BOVENGRONDSE STIKSTOFOPBRENGST MERELBEKE



BOVENGRONDSE STIKSTOFOPBRENGST OOSTENDE





De veggewassen werden ondergewerkt op 16 november 2012; er werd geen voorjaarsoogst meer uitgevoerd.





De veggewassen werden ondergewerkt op 16 november 2012; er werd geen voorjaarsoogst meer uitgevoerd.

BOVENGRONDSE STIKSTOFOPBRENGST RUKKELINGEN-LOON




De veggewassen werden geklepeld op 1 december 2012 en ondergewerkt op 12 januari 2013; er werd geen voorjaarsoogst meer uitgevoerd.




De veggewassen werden geklepeld op 1 december 2012 en ondergewerkt op 12 januari 2013; er werd geen voorjaarsoogst meer uitgevoerd.



BOVENGRONDSE KOOLSTOFOPBRENGST BOTTELARE





BOVENGRONDSE KOOLSTOFOPBRENGST LEMBERGE



BOVENGRONDSE KOOLSTOFOPBRENGST RUKKELINGEN-LOON




De veggewassen werden ondergewerkt op 15 januari 2012; er werd geen voorjaarssoogst meer uitgevoerd.






De veggewassen werden ondergewerkt op 15 januari 2012; er werd geen voorjaarssoogst meer uitgevoerd.



BOVENGRONDSE KOOLSTOFOPBRENGST SINT-DENIJS




BOVENGRONDSE KOOLSTOFOPBRENGST BOTTELARE





*2^e zaai niet geoogst eind oktober
omwille van zeer beperkte opkomst*





BOVENGRONDSE KOOLSTOFOPBRENGST MERELBEKE



BOVENGRONDSE KOOLSTOFOPBRENGST OOSTENDE





De vanggewassen werden ondergewerkt op 16 november 2012; er werd geen voorjaarsoogst meer uitgevoerd.





De vanggewassen werden ondergewerkt op 16 november 2012; er werd geen voorjaarsoogst meer uitgevoerd.

BOVENGRONDSE KOOLSTOFOPBRENGST RUKKELINGEN-LOON



De vanggewassen werden geklepeld op 1 december 2012 en ondergewerkt op 12 januari 2013; er werd geen voorjaarsoogst meer uitgevoerd.








De vanggewassen werden geklepeld op 1 december 2012 en ondergewerkt op 12 januari 2013; er werd geen voorjaarsoogst meer uitgevoerd.

MINERALE BODEMSTIKSTOF BOTTELARE


**Bottelare
HOAGENT**

Lemig zand / licht zandleem
1^e zaai: °10/08/2011
2^e zaai: °29/08/2011



*Noot: de 2 zaaitijdstoppen werden aangelegd op
2 verschillende percelen*



MINERALE BODEMSTIKSTOF BOTTELARE








MINERALE BODEMSTIKSTOF BOTTELARE




MINERALE BODEMSTIKSTOF LEMBERGE

Lemberge
ILVO



Zandleem
1^e zaai: °04/08/2011
2^e zaai: °31/08/2011




MINERALE BODEMSTIKSTOF LEMBERGE



MINERALE BODEMSTIKSTOF LEMBERGE





MINERALE BODEMSTIKSTOF RUKKELINGEN-LOON




Rukkelingen-Loon
BDB


Zwaar leem
1^e zaai: °18/08/2011
2^e zaai: °31/08/2011





Er werd geen aanvullend staal genomen eind augustus.



MINERALE BODEMSTIKSTOF RUKKELINGEN-LOON








MINERALE BODEMSTIKSTOF RUKKELINGEN-LOON




MINERALE BODEMSTIKSTOF SINT-DENIJS

Sint-Denijs
INAGRO



Licht leem
1^e zaai: °12/08/2011
2^e zaai: °30/08/2011



MINERALE BODEMSTIKSTOF SINT-DENIJS








MINERALE BODEMSTIKSTOF SINT-DENIJS




MINERALE BODEMSTIKSTOF BOTTELARE

**Bottelare
HOGENT**

Lemig zand
1^e zaai: °22/08/2012
2^e zaai: °08/09/2012








MINERALE BODEMSTIKSTOF BOTTELARE




MINERALE BODEMSTIKSTOF MERELBEKE

Merelbeke
ILVO


Lemig zand
1^e zaai: °01/08/2012
2^e zaai: °30/08/2012



MINERALE BODEMSTIKSTOF MERELBEKE








MINERALE BODEMSTIKSTOF MERELBEKE




MINERALE BODEMSTIKSTOF OOSTENDE

Oostende
INAGRO



Klei
1^e zaai: °22/08/2012
2^e zaai: °11/09/2012



MINERALE BODEMSTIKSTOF OOSTENDE








MINERALE BODEMSTIKSTOF OOSTENDE




MINERALE BODEMSTIKSTOF RUKKELINGEN-LOON

Rukkelingen-Loon
BDB

Licht leem
1^e zaai: °21/08/2012
2^e zaai: °05/09/2012




MINERALE BODEMSTIKSTOF RUKKELINGEN-LOON




MINERALE BODEMSTIKSTOF RUKKELINGEN-LOON




PROFIELEN MINERALE BODEMSTIKSTOF BOTTELARE 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF BOTTELARE 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF BOTTELARE 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF BOTTELARE 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF LEMBERGE 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF LEMBERGE 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF LEMBERGE 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF LEMBERGE 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF LEMBERGE 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF RUKKELINGEN-LOON 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF RUKKELINGEN-LOON 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF RUKKELINGEN-LOON 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF RUKKELINGEN-LOON 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF RUKKELINGEN-LOON 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF SINT-DENIJS 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF SINT-DENIJS 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF SINT-DENIJS 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF SINT-DENIJS 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF SINT-DENIJS 2011-2012




PROFIELEN MINERALE BODEMSTIKSTOF BOTTELARE




PROFIELEN MINERALE BODEMSTIKSTOF BOTTELARE




PROFIELEN MINERALE BODEMSTIKSTOF BOTTELARE




PROFIELEN MINERALE BODEMSTIKSTOF BOTTELARE




PROFIELEN MINERALE BODEMSTIKSTOF BOTTELARE




PROFIELEN MINERALE BODEMSTIKSTOF MERELBEKE




PROFIELEN MINERALE BODEMSTIKSTOF MERELBEKE




PROFIELEN MINERALE BODEMSTIKSTOF MERELBEKE




PROFIELEN MINERALE BODEMSTIKSTOF MERELBEKE




PROFIELEN MINERALE BODEMSTIKSTOF MERELBEKE




PROFIELEN MINERALE BODEMSTIKSTOF OOSTENDE




PROFIELEN MINERALE BODEMSTIKSTOF OOSTENDE




PROFIELEN MINERALE BODEMSTIKSTOF OOSTENDE




PROFIELEN MINERALE BODEMSTIKSTOF OOSTENDE




PROFIENEN MINERALE BODEMSTIKSTOF RUKKELINGEN-LOON




PROFIENEN MINERALE BODEMSTIKSTOF RUKKELINGEN-LOON




PROFIENEN MINERALE BODEMSTIKSTOF RUKKELINGEN-LOON




PROFIENEN MINERALE BODEMSTIKSTOF RUKKELINGEN-LOON




PROFIENEN MINERALE BODEMSTIKSTOF RUKKELINGEN-LOON




Bijlage 11: Gesimuleerde en gemeten minerale N op het proefperceel te Sint-Denijs (1e proefjaar)




Bijlage 11: Gesimuleerde en gemeten minerale N op het proefperceel te Sint-Denijs (1e proefjaar)




Bijlage 11: Gesimuleerde en gemeten minerale N op het proefperceel te Sint-Denijs (1e proefjaar)




Bijlage 11: Gesimuleerde en gemeten minerale N op het proefperceel te Sint-Denijs (1e proefjaar)




Bijlage 11: Gesimuleerde en gemeten minerale N op het proefperceel te Sint-Denijs (1e proefjaar)




Bijlage 12: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Sint-Denijs (1e proefjaar)




Bijlage 12: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Sint-Denijs (1e proefjaar)




Bijlage 12: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Sint-Denijs (1e proefjaar)




Bijlage 12: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Sint-Denijs (1e proefjaar)




Bijlage 13: Gesimuleerde en gemeten minerale N op het proefperceel te Bottelare (2e proefjaar)




Bijlage 13: Gesimuleerde en gemeten minerale N op het proefperceel te Bottelare (2e proefjaar)




Bijlage 13: Gesimuleerde en gemeten minerale N op het proefperceel te Bottelare (2e proefjaar)




Bijlage 13: Gesimuleerde en gemeten minerale N op het proefperceel te Bottelare (2e proefjaar)




Bijlage 13: Gesimuleerde en gemeten minerale N op het proefperceel te Bottelare (2e proefjaar)




Bijlage 14: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Bottelare (2e proefjaar)




Bijlage 14: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Bottelare (2e proefjaar)




Bijlage 14: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Bottelare (2e proefjaar)




Bijlage 14: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Bottelare (2e proefjaar)




Bijlage 15: Gesimuleerde en gemeten minerale N op het proefperceel te Merelbeke (2e proefjaar)




Bijlage 15: Gesimuleerde en gemeten minerale N op het proefperceel te Merelbeke (2e proefjaar)




Bijlage 15: Gesimuleerde en gemeten minerale N op het proefperceel te Merelbeke (2e proefjaar)




Bijlage 15: Gesimuleerde en gemeten minerale N op het proefperceel te Merelbeke (2e proefjaar)




Bijlage 15: Gesimuleerde en gemeten minerale N op het proefperceel te Merelbeke (2e proefjaar)




Bijlage 16: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Merelbeke (2e proefjaar)




Bijlage 16: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Merelbeke (2e proefjaar)




Bijlage 16: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Merelbeke (2e proefjaar)




Bijlage 16: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Merelbeke (2e proefjaar)




Bijlage 17: Gesimuleerde en gemeten minerale N op het proefperceel te Rukkelingen-Loon (2e proefjaar)




Bijlage 17: Gesimuleerde en gemeten minerale N op het proefperceel te Rukkelingen-Loon (2e proefjaar)




Bijlage 17: Gesimuleerde en gemeten minerale N op het proefperceel te Rukkelingen-Loon (2e proefjaar)




Bijlage 17: Gesimuleerde en gemeten minerale N op het proefperceel te Rukkelingen-Loon (2e proefjaar)




Bijlage 17: Gesimuleerde en gemeten minerale N op het proefperceel te Rukkelingen-Loon (2e proefjaar)




Bijlage 18: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Rukkelingen-Loon (2e proefjaar)




Bijlage 18: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Rukkelingen-Loon (2e proefjaar)




Bijlage 18: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Rukkelingen-Loon (2e proefjaar)




Bijlage 18: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Rukkelingen-Loon (2e proefjaar)




Bijlage 19: Gesimuleerde en gemeten minerale N op het proefperceel te Bottelare (1e proefjaar)




Bijlage 19: Gesimuleerde en gemeten minerale N op het proefperceel te Bottelare (1e proefjaar)




Bijlage 19: Gesimuleerde en gemeten minerale N op het proefperceel te Bottelare (1e proefjaar)




Bijlage 19: Gesimuleerde en gemeten minerale N op het proefperceel te Bottelare (1e proefjaar)




Bijlage 20: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Bottelare (1e proefjaar)



Bijlage 20: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Bottelare (1e proefjaar)



Bijlage 20: Gesimuleerde en gemeten bovengrondse N-opbrengsten op het proefperceel te Bottelare (1e proefjaar)



Bijlage 21: Analyse van de residuen

TABEL 1: GEMIDDELDE AFWIJING TUSSEN GESIMULEERDE N_{MIN}-GEHALTEN EN METINGEN OP DE PROEFPERCELEN (IN KG N HA⁻¹).

Proefjaar Locatie Zaaitijdstip	2011-2012						2012-2013						gemiddeld	
	LEM		STD		BOT		MER		RUK		BOT			
	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2
BR0	-7,4	7,0	-6,7	-6,7	6,2	-10,7	7,2	7,2	-3,5	-3,5	-9,4	-9,4	-2,3	-2,7
BR60	-8,6	6,1	8,2	8,2	17,8	-6,8	19,6	19,6	9,7	9,7	-2,3	-2,3	7,4	5,8
BR120	-11,5	-5,6	16,6	16,6	18,9	-21,5	10,0	10,0	38,2	38,2	0,7	0,7	12,1	6,4
BR	-9,2	2,5	6,0	6,0	14,3	-13,0	12,3	12,3	14,8	14,8	-3,6	-3,6	5,8	3,2
GM0	-4,2	0,7	-1,1	5,3	-7,7	-18,1	4,8	2,8	4,7	-12,5	-3,9	-36,6	-1,2	-9,7
GM60	-5,4	0,1	-1,2	2,4	-7,1	-28,3	10,2	14,1	10,0	-8,8	0,6	-52,3	1,2	-12,1
GM120	-4,2	-8,6	-0,6	1,9	-3,4	-25,9	11,6	14,6	30,1	24,8	-1,0	-61,3	5,4	-9,1
GM	-4,6	-2,6	-1,0	3,2	-6,1	-24,1	8,9	10,5	14,9	1,2	-1,4	-50,1	1,8	-10,3
IRO	-4,5	4,5	9,3	9,9	-2,0	-2,2	3,8	-11,8	14,6	-1,2	-0,5	-31,0	3,4	-5,3
IR60	-7,8	-0,8	2,4	2,7	-7,3	-7,1	0,4	-12,5	16,9	-3,4	-4,2	-44,8	0,1	-11,0
IR120	-12,2	-0,7	-0,3	0,1	-13,1	-17,1	1,7	-30,0	19,1	20,7	-3,9	-56,1	-1,5	-13,9
IR	-8,2	1,0	3,8	4,2	-7,5	-8,8	1,9	-18,1	16,9	5,4	-2,9	-43,9	0,7	-10,0
JH0	-2,5	-0,4	12,9	14,9	8,5	14,2	1,7	11,9	14,7	-2,3	4,6	-16,3	6,6	3,7
JH60	-2,7	-1,0	9,9	18,8	14,7	20,4	2,7	29,2	29,5	6,8	13,5	-23,6	11,3	8,4
JH120	-9,3	-11,5	12,1	32,9	21,9	21,1	1,6	17,1	63,7	47,9	24,0	-25,5	19,0	13,7
JH	-4,8	-4,3	11,7	22,2	15,0	18,6	2,0	19,4	35,9	17,5	14,0	-21,8	12,3	8,6
GK0 + fixatie	-3,0	0,3	4,7	1,7	-	-	12,4	3,5	26,0	7,8	4,6	-27,5	8,9	-2,8
GK60 + fixatie	-3,5	-2,0	-1,5	2,4	-	-	15,0	15,3	48,8	12,8	7,8	-26,8	13,3	0,4
GK120 + fixatie	-8,0	-9,3	9,5	17,1	-	-	30,1	14,2	77,5	51,1	15,0	-39,1	24,8	6,8
GK + fixatie	-4,8	-3,6	4,2	7,1	-	-	19,2	11,0	50,7	23,9	9,1	-31,1	15,7	1,4
GK0 - fixatie	1,3	4,4	13,3	14,9	-	-	7,8	3,8	27,0	7,3	5,7	-26,0	11,0	0,9
GK60 - fixatie	-1,8	-0,4	-0,8	6,3	-	-	10,2	9,0	47,0	11,2	3,5	-29,0	11,6	-0,6
GK120 - fixatie	-7,3	-9,1	2,3	13,6	-	-	21,6	4,7	71,1	48,3	9,3	-42,9	19,4	2,9
GK - fixatie	-2,6	-1,7	4,9	11,6	-	-	13,2	5,8	48,4	22,3	6,2	-32,6	14,0	1,1
gemiddeld	-5,7	-1,5	4,9	9,1	4,0	-6,8	9,6	6,8	30,3	14,2	3,6	-30,5	7,8	-1,5
gemiddeld Z1+Z2	-3,6		7,0		-1,4		8,2		22,2		-13,5		3,2	

Bijlage 21: Analyse van de residuen

TABEL 2: GEMIDDELTE ABSOLUTE AFWIJING TUSSEN GESIMULEERDE N_{MIN}-GEHALTEN EN METINGEN OP DE PROEFPERCELEN (IN KG N HA⁻¹).

Proefjaar Locatie Zaaitijdstip	2011-2012						2012-2013						gemiddeld	
	LEM		STD		BOT		MER		RUK		BOT			
	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2
BR0	8,3	9,9	7,6	7,6	10,3	14,4	9,7	9,7	14,4	14,4	14,6	14,6	10,8	11,7
BR60	8,9	6,4	8,2	8,2	20,9	11,3	27,8	27,8	11,2	11,2	11,0	11,0	14,7	12,6
BR120	11,5	11,1	20,4	20,4	19,0	27,0	29,7	29,7	38,2	38,2	16,8	16,8	22,6	23,9
BR	9,6	9,1	12,1	12,1	16,7	17,6	22,4	22,4	21,2	21,2	14,1	14,1	16,0	16,1
GM0	5,2	3,7	8,9	5,4	12,1	18,1	6,3	9,4	13,3	15,3	9,1	40,1	9,2	15,3
GM60	5,5	5,3	7,3	3,8	18,0	28,3	10,2	16,5	11,7	9,7	19,2	62,0	12,0	20,9
GM120	5,6	11,8	6,8	4,0	25,8	39,4	13,5	29,8	30,1	24,8	15,0	61,4	16,1	28,5
GM	5,4	7,0	7,7	4,4	18,6	28,6	10,0	18,6	18,4	16,6	14,4	54,5	12,4	21,6
IRO	5,8	4,7	9,9	9,9	12,6	6,8	4,7	13,1	18,5	24,3	9,2	33,3	10,1	15,3
IR60	7,8	3,9	6,9	5,1	15,4	9,1	2,8	16,1	21,1	19,6	12,4	44,8	11,0	16,5
IR120	12,2	8,4	7,5	4,0	15,0	17,1	4,4	31,2	21,6	21,4	14,2	56,6	12,5	23,1
IR	8,6	5,7	8,1	6,4	14,3	11,0	4,0	20,1	20,4	21,8	11,9	44,9	11,2	18,3
JH0	3,9	11,1	12,9	14,9	16,6	26,0	9,0	13,2	15,2	21,6	18,3	24,4	12,6	18,5
JH60	4,5	11,4	9,9	18,8	23,0	34,4	7,0	30,5	33,0	18,8	30,3	31,6	17,9	24,3
JH120	10,9	19,3	12,1	32,9	26,6	38,8	11,5	22,5	63,7	47,9	35,4	28,3	26,7	31,6
JH	6,4	13,9	11,7	22,2	22,1	33,1	9,1	22,1	37,3	29,4	28,0	28,1	19,1	24,8
GK0 + fixatie	3,5	7,3	11,5	13,0	-	-	16,1	6,5	37,8	25,3	9,5	31,9	15,7	16,8
GK60 + fixatie	3,9	6,1	14,6	16,7	-	-	20,5	21,7	48,8	27,7	17,7	31,0	21,1	20,6
GK120 + fixatie	8,0	9,6	16,8	24,9	-	-	34,6	21,2	77,5	51,1	22,3	43,6	31,8	30,1
GK + fixatie	5,1	7,7	14,3	18,2	-	-	23,7	16,5	54,7	34,7	16,5	35,5	22,9	22,5
GK0 - fixatie	4,5	7,1	13,7	20,0	-	-	11,5	10,3	39,4	24,9	13,4	30,6	16,5	18,6
GK60 - fixatie	5,2	3,4	11,9	16,6	-	-	15,4	17,1	47,0	26,5	19,5	32,2	19,8	19,2
GK120 - fixatie	7,3	9,2	18,3	24,0	-	-	25,9	18,3	71,1	48,3	21,2	46,5	28,8	29,3
GK - fixatie	5,6	6,6	14,6	20,2	-	-	17,6	15,3	52,5	33,3	18,0	36,4	21,7	22,3
gemiddeld	6,8	8,3	11,4	13,9	18,0	22,6	14,5	19,2	34,1	26,2	17,2	35,6	17,0	20,9
gemiddeld Z1+Z2	7,6		12,7		20,3		16,8		30,1		26,4		19,0	

Bijlage 21: Analyse van de residuen

TABEL 3: GEMIDDELTE AFWIJING TUSSEN GESIMULEERDE BOVENGRONDSE N-OPBRENGSTEN EN METINGEN OP DE PROEFPERCELEN (IN KG N HA⁻¹).

Proefjaar Locatie Zaaitijdstip	2011-2012						2012-2013						gemiddeld	
	LEM		STD		BOT		MER		RUK		BOT			
	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2
GM0	-0,6	5,9	-6,6	-4,5	25,4	20,5	4,9	7,7	9,5	12,6	18,8	30,9	8,6	12,2
GM60	-1,9	2,4	4,6	3,0	21,5	22,0	21,6	18,4	27,6	20,2	21,7	36,3	15,9	17,0
GM120	-9,9	-0,8	18,3	13,9	33,8	35,6	21,0	3,2	47,2	22,9	30,4	49,9	23,5	20,8
GM	-4,1	2,5	5,4	4,1	26,9	26,0	15,8	9,7	28,1	18,6	23,7	39,0	16,0	16,7
IRO	3,8	10,4	10,0	13,0	36,3	34,0	28,0	20,1	8,2	7,3	24,8	31,3	18,5	19,4
IR60	-11,9	-3,2	19,1	21,8	51,2	42,1	31,7	43,9	17,5	19,2	34,8	46,7	23,7	28,4
IR120	-12,3	3,1	34,4	38,5	52,9	50,3	19,8	39,2	29,0	21,1	34,1	66,6	26,3	36,5
IR	-6,8	3,5	21,1	24,4	46,8	42,2	26,5	34,4	18,3	15,9	31,2	48,2	22,9	28,1
JH0	-3,5	11,2	-7,7	2,9	40,6	-5,0	39,4	2,8	12,3	4,2	24,8	23,2	17,7	6,5
JH60	-8,7	9,2	2,4	5,9	39,4	-14,8	56,9	-0,6	0,3	4,7	27,6	23,8	19,6	4,7
JH120	-8,6	-0,8	26,9	-5,8	31,2	-22,9	72,5	-25,0	-4,0	0,6	34,4	13,5	25,4	-6,7
JH	-6,9	6,5	7,2	1,0	37,1	-14,2	56,3	-7,6	2,9	3,1	28,9	20,1	20,9	1,5
GK0 + fixatie	5,0	11,9	23,7	21,2	-	-	10,2	5,9	7,6	-	8,8	6,5	11,1	11,4
GK60 + fixatie	-5,2	-2,0	13,0	12,2	-	-	31,3	23,4	-8,3	-	-9,1	8,9	4,4	10,6
GK120 + fixatie	-8,3	-7,9	-4,5	1,8	-	-	44,7	30,9	-13,7	-	-24,7	5,2	-1,3	7,5
GK + fixatie	-2,8	0,7	10,7	11,7	-	-	28,7	20,1	-4,8	-	-8,3	6,9	4,7	9,8
GK0 - fixatie	-0,7	14,5	10,8	10,2	-	-	9,2	8,5	6,3	-	7,2	7,4	6,6	10,1
GK60 - fixatie	-8,3	6,6	25,4	25,7	-	-	2,8	18,1	-0,3	-	8,1	18,6	5,6	17,3
GK120 - fixatie	-8,8	4,8	25,9	24,0	-	-	-33,6	12,1	-0,1	-	3,6	19,2	-2,6	15,0
GK - fixatie	-5,9	8,6	20,7	20,0	-	-	-7,2	12,9	2,0	-	6,3	15,1	3,2	14,1
gemiddeld	-5,3	4,4	13,0	12,2	36,9	18,0	24,0	13,9	9,3	12,5	16,4	25,9	15,7	14,5
gemiddeld Z1+Z2	-0,5		12,6		27,4		19,0		10,9		21,1		15,1	

Bijlage 21: Analyse van de residuen


TABEL 4: GEMIDDELDE ABSOLUTE AFWIJING TUSSEN GESIMULEERDE BOVENGRONDSE N-OPBRENGSTEN EN METINGEN OP DE PROEFPERCELEN (IN KG N HA⁻¹).

Proefjaar Locatie Zaaitijdstip	2011-2012						2012-2013						gemiddeld	
	LEM		STD		BOT		MER		RUK		BOT			
	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2	zaai1	zaai2
GM0	9,1	5,9	9,6	5,3	25,4	20,5	7,4	7,7	11,2	12,6	18,8	30,9	13,6	13,8
GM60	4,0	5,3	8,9	3,0	21,5	22,0	21,6	18,4	27,6	20,2	21,7	36,3	17,6	17,5
GM120	9,9	7,8	18,3	13,9	33,8	35,6	21,0	19,6	47,2	22,9	30,4	49,9	26,8	24,9
GM	7,6	6,3	12,3	7,4	26,9	26,0	16,7	15,2	28,7	18,6	23,7	39,0	19,3	18,8
IRO	3,8	10,4	10,0	13,0	36,3	34,0	28,0	20,1	8,2	7,3	24,8	31,3	18,5	19,4
IR60	11,9	7,7	19,1	21,8	51,2	42,1	31,7	43,9	17,5	19,2	34,8	46,7	27,7	30,2
IR120	12,3	6,7	34,4	38,5	52,9	50,3	19,8	39,2	29,0	21,1	34,1	66,6	30,4	37,1
IR	9,3	8,3	21,1	24,4	46,8	42,2	26,5	34,4	18,3	15,9	31,2	48,2	25,5	28,9
JH0	4,7	11,2	18,6	3,5	40,6	5,6	39,4	13,3	14,0	4,2	24,8	23,2	23,7	10,1
JH60	8,7	9,2	10,1	16,4	39,4	14,8	56,9	12,0	19,7	5,5	27,6	23,8	27,0	13,6
JH120	9,7	1,4	26,9	18,3	34,0	22,9	72,5	25,0	22,6	4,8	34,4	17,3	33,4	14,9
JH	7,7	7,3	18,5	12,7	38,0	14,4	56,3	16,7	18,8	4,8	28,9	21,4	28,0	12,9
GK0 + fixatie	6,0	11,9	23,7	21,2	-	-	10,2	5,9	7,6	-	11,8	6,5	11,9	11,4
GK60 + fixatie	5,2	3,0	15,3	12,2	-	-	31,3	23,4	8,3	-	13,4	8,9	14,7	11,9
GK120 + fixatie	8,3	7,9	12,5	9,6	-	-	44,7	30,9	13,7	-	24,7	5,2	20,8	13,4
GK + fixatie	6,5	7,6	17,2	14,4	-	-	28,7	20,1	9,9	-	16,6	6,9	15,8	12,2
GK0 - fixatie	6,2	14,5	11,0	10,2	-	-	9,2	8,5	6,3	-	8,1	7,4	8,2	10,1
GK60 - fixatie	8,3	6,6	25,4	25,7	-	-	12,3	18,1	5,3	-	16,3	18,6	13,5	17,3
GK120 - fixatie	8,8	6,7	25,9	24,0	-	-	33,6	18,9	0,7	-	15,8	19,2	17,0	17,2
GK - fixatie	7,8	9,2	20,8	20,0	-	-	18,3	15,2	4,1	-	13,4	15,1	12,9	14,9
gemiddeld	7,8	7,7	18,0	15,8	37,2	27,5	29,3	20,3	15,9	13,1	22,8	26,1	21,8	18,4
gemiddeld Z1+Z2	7,8		16,9		32,4		24,8		14,5		24,4		20,1	

BRIEF DESCRIPTION OF THE EU-ROTATE_N MODEL

8/08/2007

EU-Rotate_N



EU-Rotate_N is a four year project, funded by the European Commission within the Fifth Framework Programme, and is co-ordinated by Dr Clive Rahn at Warwick HRI. Partner organizations include HDRA UK, IGZ and BOLAP Germany , IVIA and CSIC Spain, Bioforsk Norway, DJF Denmark and ISOR Italy.

The developers can be contacted eurotaten@warwick.ac.uk or visit the Website at www.warwick.ac.uk/go/eurotaten

The Integrated Model - description of the model sub-routines.

Rahn CR, Zhang K, Lillywhite R, Ramos C, Doltra J, de Paz J M, Riley H, Fink M, Nendel C, Thorup Kristensen K, Piro F, Venezia A, Firth C, Schmutz U, Raynes F, Strohmeyer K.

Where the model sub-routines are based on existing models or existing published algorithms these are referenced to the original source. Where new sub routines have been added which are based on new science such as the root routines these are described in more detail.

1 Model structure and timestep.....	2
2 Description of the soil.....	4
3 The water module	5
4 Mineralisation Module.....	6
5 Snow and Frost Module.....	7
6 Root Module	8
7 N uptake	10
8 Crop growth and critical N	11
9 Fertility Building Crops	12
10 Economics Module	13

1 Model structure and timestep

The model consists of a number of subroutines to simulate the growth both below and above ground, nitrogen mineralisation from the soil and crop residues, subsequent N uptake and balance between supply and demand to regulate growth. These will all be regulated by weather factors such as rainfall, temperature and radiation. Routines simulate the flow of water and nitrogen into the plant, subsequent evapotranspiration or leaching.

The sub-routines operate in the following order, utilising data from soil properties, residues, fertiliser and weather data where appropriate.

1. The soil N mineralization: calculates soil N mineralization in the top 30cm soil depth from soil organic matter, crops residues and organic and inorganic N fertilisers;
2. The potential maximum increment in shoot dry weight: calculated on the assumption of no restriction from N-deficiency and water stress;
3. The potential maximum N-uptake: calculated from the product of potential maximum dry weight and the critical %N for a crop of that size;
4. The root distribution: calculates the rooting depth and width, and root length distribution in the root zone;

5. The actual N uptake: calculates the amount of N that the roots can take from the root zone;
6. The actual %N in the plant: calculated from the N uptake, the amount of N in the plant on the previous day, and the dry weight of the plant calculated for the previous day;
7. The snow dynamics and frost depth: calculates the snow depth, depths of frost and of any thawed layer above the frost layer, actual infiltration water for winter climate;
8. The surface runoff: calculates the surface runoff caused by heavy rain;
9. The potential transpiration and evaporation rate: calculated from the reference transpiration and a crop coefficient varying with crop development using a FAO approach;
10. The water drainage and redistribution: calculates soil water distribution in the entire soil domain from infiltration and soil properties;
11. The water removal: calculates the amounts of water that the roots can extract from the root zone and via evaporation;
12. The actual shoot dry weight increment: calculated from its current dry weight, the calculated %N and reduction in transpiration.
13. At harvest the amount of marketable crop and its gross margin is calculated using the Economics module.

Modules 1-12 are daily routines. Whether modules 7 & 8 are called is dependent on the information in the input file. During growth modules 1-12 are called on a daily basis, while modules 1, 7-11 are called when the model simulates fallow crops. At harvest module 13 is also called.




Figure 1: The organisation of the main model sub_modules

2 Description of the soil

In the model the soil is divided into 40 5 cm thick vertical layers. After planting these layers are split horizontally into 5cm wide cells.. The number of cells horizontally depends on row width. As soon as the crop is harvested OR when the residues are incorporated the horizontal elements are merged into one unit until the next crop is planted. The description of the soil in this way allows for more accurate simulation of root growth than was possible in the original N_ABLE model. While the crop is growing all the processes described below are all simulated at the element level.

The properties of the soil layers are provided by the user of the model and include the water content at permanent wilting point, field capacity and at the saturated level. These control water availability to the plant and allow calculation of drainage. Mineralisation and losses of nitrogen by denitrification is adjusted for water content. Other inputs include pH which allows for the simulation of N losses where urea fertilisers are used. The amount of organic matter levels affects the supply of N from mineralisation. Clay and sand contents are used to calculate urea solution and hydrolysis, ammonia volatilisation from top layer, decay rate coefficients, denitrification.

3 The water module

The water module has different parts that calculate basically the following:

- Crop evapotranspiration (soil evaporation and transpiration)
- Effective water infiltration (applied water minus runoff)
- Drainage
- Water redistribution in soil

Crop evapotranspiration is calculated using basically the FAO approach (Allen et al., 1998). The main parameters that enter in these calculations are those related to the evaporative demand of the atmosphere, summarized by the reference evapotranspiration (ET_o) and a crop coefficient that varies with crop development.

The effects of water stress on plant growth are considered assuming that the reduction in dry matter accumulation due to water deficit is proportional to the transpiration reduction (Hanks, 1983; Shani and Dudley, 2001).

Water infiltration and redistribution in soil follows a capacitance approach, similar to the one in the N_ABLE model but that has been modified using a drainage coefficient that allows the water transfer between layers above field capacity to be done progressively (in more than one day) and more or less rapidly depending on soil type (Ritchie, 1998). Drainage at any depth is given as the water downward flow of the cell elements at this depth. The module also accounts for upward/downward capillary flow by adopting a soil water normalised diffusion approach (Rose, 1968 and Ritchie, 1998). The main parameters that define the hydraulic soil properties such as the water content at field capacity and wilting point, are input by the user for the different soil layers, although default values depending on soil texture are available.

Runoff is calculated using the approach by the U.S. National Resource Conservation Service (NRCS, formerly the Soil Conservation Service) based on studies of small (< 800 ha) agricultural watersheds across the United States.

References

- Allen R.G., Pereira L.S., Raes D., Smith M.(1998). Crop evapotranspiration. Guidelines for computing crop water requirements. FAO Irrigation and Drainage Paper n° 56, FAO, Roma.
- Haan, C.T., B.J. Barfield and J.C. Hayes (1994) Design hydrology and sedimentation for small catchments, Academic Press, New York, 588p.
- Hanks R.J. 1983. Yield and water-use relationships: An overview. In H.M. Taylor et al. (ed.) Limitations to efficient water use in crop production. ASA, Madison, WI., pp. 393-411.
- Ritchie, J.T. (1998). Soil water balance and plant water stress. G.Y. Tsuji et al. (eds): Understanding options for agricultural production p. 41-54.
- Rose, D. (1968). Water movement in porous materials III. Evaporation of water from soil. Brit. J. Appl. Phys. ser 2(1): 1779-1791.
- Shani U. and L. M. Dudley. 2001. Field studies of crop response to water and salt stress. Soil Sci. Soc. Am. J. 65:1522–1528.

4 Mineralisation Module

Nitrogen (N) release from soil organic matter and from different kinds of N containing materials added to the soil for fertilising is calculated in a module that hosts the following routines:

- N mineralisation from both soil bound and freshly added organic matter
- N volatilisation from added manures and slurries
- Hydrolysis of urea and volatilisation of NH_3

The concept of N mineralisation from organic matter is based on the routines used in the DAISY model (Hansen et al., 1990). Three pairs of conceptual pools (soil organic matter, soil microbial biomass added organic matter), each representing a rapidly decomposable and a slowly decomposable class of N containing organic substances, respectively, describe C dynamics in the soil. Decomposition rate coefficients are temperature and moisture dependent and reflect the environmental conditions of the simulated site. Decay and maintenance rates of soil microbial biomass are additionally influenced by soil clay content. Efficiency parameters determine the loss of CO_2 during the single turn-over processes. N release as NH_4^+ is a consequence of C lost as CO_2 from the system that maintains fixed C to N ratios in the different pools. Processes of nitrification and denitrification are implemented to complete the turn-over model.

Residues of crops simulated with the crop growth model enter the mineralisation routine with a dynamic C to N ratio, which reflects the growth conditions of the crop during season with respect to N supply. The variable C to N ratio is assigned to the rapidly decomposing part of the material, while the remaining part is considered to decompose slowly, having a fixed C to N ratio. Decomposition rate coefficients of both pools are also fixed (Abrahamsen, 2000). C to N ratios and partitioning coefficients for crop residues are derived from stepwise chemical digestion experiments (Jensen et al. 2005). Manure and slurry properties are taken from DAISY parameterisations (Abrahamsen, 2000).

N volatilisation from soil applied manures and slurries are described using an empirical relation implemented in the ALFAM model (Søgaard et al., 2002). A soil pH dependency factor was introduced by fitting data from He et al. (1999) to Michaelis-Menten kinetics and subsequently normalising the relation between pH and volatilisation half life time to pH 7.0.

Hydrolysis of and gaseous N loss from applied urea fertiliser is calculated based on routines of the AMOVOL model (Sadeghi et al., 1988), taking into account the temperature dependent equilibrium between ammonium ions, solved and gaseous ammonia as well as the effect of soil organic matter, soil temperature and soil water potential on the hydrolysis process itself. An atmospheric resistance parameter finally governs the loss of gaseous ammonia from the top soil.

Literature

Abrahamsen, P., Hansen, S., 2000. Daisy: an open soil-crop-atmosphere system model. *Environmental Modelling & Software* 15, 313-330.

Hansen, S., Jensen, H.E., Nielsen, N.E., Svendsen, H., 1990. Daisy - A Soil Plant Atmosphere System Model. NPO Research from the National Agency of Environmental Protection No. A 10. 272 pp.

He, Z.L., A.K. Alva, D.V. Calvert und D.J. Banks (1999): Ammonia volatilization from different fertilizer sources and effects of temperature and soil pH. *Soil Science*, 164 (10): 750-758.

Jensen, L.S., Salo, T., Palmason, F., Breland, T.A., Henriksen, T.M., Stenberg, B., Pedersen, A., Lundström, C., Esala, M., 2005. Influence of biochemical quality on C and N mineralisation from a broad variety of plant materials in soil. *Plant and Soil* 273, 307-326.

Sadeghi, A.M., K.J. McInnes, D.E. Kissel, M.L. Cabrera, J.K. Koelliker und E.T. Kanemasu (1988): Mechanistic model for predicting ammonia volatilization from urea. In: B.R. Bock und D.E. Kissel (Hrsg.), *Ammonia volatilization from urea fertilizers*. National Fertilizer Development Centre, Tennessee Valley Authority, Muscle Shoals, Alabama, 67- 92.

Søgaard, H.T., S.G. Sommer, N.J. Hutchings, J.F.M. Huijsmans, D.W. Bussink und F. Nicholson (2002): Ammonia volatilization from field-applied animal slurry - the ALFAM model. *Atmospheric Environment*, 36 (20): 3309-3319

5 Snow and Frost Module

Routines have been developed that allow the calculation of snow depth and density, water storage in snow and water melting from the snow pack, using daily input of air temperature.

The original snow model was developed at the University of Helsinki by Vehviläinen & Lohvansuu (1991) for calculating water equivalent, but modified by Tuomo Karvonen (see internet reference) to calculate snow depth, which is important for determining soil freezing and thawing. We have further modified this model and have calibrated it by iterative simulation using a 10-year dataset from Norway, as described by Riley and Bonesmo (2005). The approach has later been validated with independent data.

The chosen soil frost model is based on two approaches, one for freezing and one for thawing. The approach for soil freezing was proposed by Olsen and Haugen (1997), at the Norwegian University of Life Sciences, Ås, assuming uniform thermal properties throughout the profile. Values for the latter properties are taken from the Swedish SOIL model (Jansson, 1991). The model requires input of surface temperature as modified by the snow pack. The approach used for thawing is that in the ECOMAG model developed at the University of Oslo (Molotov et al., 1999). We have validated both freezing and thawing processes for Norwegian conditions.

The snow and frost calculation procedures, including all parameters used, are described in detail in a programming note (Riley, 2004a). This note also describes how these processes interact with water infiltration and associated processes such as leaching. In brief, it is assumed that infiltration ceases when soil freezes. During snowmelt and soil thaw, an amount of water equal to the difference between field capacity and total porosity is stored for later infiltration, whilst the remainder passes

to surface runoff. An example of the interaction of frost with runoff appeared in an article in the second project newsletter (Riley, 1994b).

Literature cited

Jansson, P.E. 1991. Simulation modell for soil water and heat conditions. Report 165, Swedish University of Agricultural Sciences, Uppsala, 72 pp.

Karvonen T. (internet publication). Model of snow accumulation and snowmelt.
www.water.hut.fi/wr/kurssit/Yhd-12.135/kirja/paa_e.htm

Motovilov, Y.G., L. Gottschalk, K. Engeland, & A. Belokurov, 1999. ECOMAG: Regional model of hydrological cycle. Application to the NOPEX region. Dept. Geophysics, University of Oslo, Report no. 1051, 88 pp.

Olsen, P.A. & L.E. Haugen, 1997. Jordas termiske egenskaper. Dept. Soil and Water Science, Agricultural University of Norway, Rapport nr. 8, 14 pp. (in Norwegian).

Riley, H. 2004a. Description of submodels on snow dynamics and frost depth. Programming note of 18th June 2004, 9 pp.

Riley, H. 2004b. New submodel: Snow and Frost Article intended in newsletter in June 2004,

Riley, H. and Bonesmo H. 2005. Modelling of snow and freeze-thaw cycles in the EU-rotate_N decision support system. *Planteforsk Grønn Kunnskap (e)* vol. 9, no. 112, 8 pp.

Vehviläinen, B. & Lohvansuu, J. 1991. The effects of climate on water discharges and snow cover in Finland. *Hydrological Sciences* 36 (2): 109-121.

6 Root Module

The root model calculation consists of three main parts: 1) first the physical extension of the root system is calculated, 2) then the total root length of the crop is calculated, and 3) finally the distribution of the root system depending on depth and distance from the crop row is calculated.

The depth development of the root system is calculated from the accumulated temperature sum from crop planting. After a lag period (ddglag) the rooting depth increases linearly with temperature sum. The length of the lag period and the rate of rooting depth development are controlled with crop specific parameter values. A crop specific base temperature (Tmin) for calculation of root growth is also used, and Tmax is then set to Tmin+20 °C. This approach to simulation of crop rooting depth is based on a number of studies showing good linear relationships between accumulated temperature sum and rooting depth (Kristensen & Thorup-Kristensen, 2004; Thorup-Kristensen, 2001; Thorup-Kristensen, 2006; Thorup-Kristensen, 1998; Thorup-Kristensen & Van den Boogaard, 1998; Thorup-Kristensen & Van den Boogaard, 1999; Kage et al., 2000).

$$rz=zstart+((cumuT-ddglag)*Krz) \quad (1)$$

Horizontal root extension is calculated in the same way, but for each soil layer the calculation starts when the roots reach this layer rather than when the crop is planted. In this way horizontal root growth starts progressively later at larger depths.

$$rx=xstart+((cumuT-ddglag)*Krx) \quad (2)$$

Crop root length is then calculated as a function of 1) crop biomass, 2) crop growth stage, and 3) the parameter value of root class "Rtclss". The root biomass is calculated as a function of aboveground crop biomass, a fraction which declines with crop size, but increases with Rtclss ($1 < 2 < 3$), to allow for crops with different root/shoot ratios. Total root length is then calculated from the simulated root biomass and a fixed specific root length which is used for all crops.

Root length is distributed spatially into a 2D array of soil units. Root models in crop/soil simulation models are mostly 1D; i.e. root density varies only with depth. However, most vegetable crops are grown as row crops, and the 2D approach was used to be able to simulate the effects of the row crop structure on crop rooting and uptake of water and nitrogen. Root distribution is calculated to a maximum depth of 2 m, and to a max width of half crop row distance. The soil units used in this array are 0.05 by 0.05 m.

The root length declines by a logarithmic function from the topsoil downwards, as originally proposed by Gerwitz and Page (1974), and from the crop row to the interrow soil. However, contrary to Gerwitz and Page (1974) we include a value for rooting depth, under which root density declines fast to zero. In the modified forms of the Gerwitz and Page equation used in other models, a rooting depth has also been included, but in these approaches the root density at maximum rooting depth has been constant, meaning that subsoil root density will always be low, and variation in root length will practically only be found in the topsoil. With our approach we allow higher root densities in the subsoil, but compared to the original equation from Gerwitz and Page (1974) our setup allows relatively high root density at the simulated rooting depth without leading to significant root density also in layers below that. In our approach the steepness of the logarithmic decline is controlled by one parameter for the vertical distribution (az) and another parameter for the horizontal distribution (ax).

$$\text{rootlength}(i,j)=(lz(i)*\exp(-ax*x)) \quad (3)$$

Ideally, the model should have been a 3D model, to simulate also the effect of plant distance within the crop row. At the early growth stages where root width is less than plant to plant distance within the row, an arbitrary function is used to reduce N uptake capacity accordingly.

Reference List

GERWITZ A, PAGE E.R. 1974 Empirical mathematical-model to describe plant root systems. J. Appl. Ecol. 11: 773-781

- KAGE,H., KOCHLER,M. & STUTZEL,H. 2000. Root growth of cauliflower (*Brassica oleracea* L. botrytis) under unstressed conditions: measurement and modelling. *Plant & Soil* 223, 131-145.
- KRISTENSEN,H.L. & THORUP-KRISTENSEN,K. 2004. Uptake of ¹⁵N labeled nitrate by root systems of sweet corn, carrot and white cabbage from 0.2 to 2.5 m depth. *Plant & Soil* 265, 93-100.
- THORUP-KRISTENSEN,K. 1998. Root Growth of Green Pea (*Pisum sativum* L.) Genotypes. *Crop Sci.* 38, 1445-1451.
- THORUP-KRISTENSEN,K. 2001. Are differences in root growth of nitrogen catch crops important for their ability to reduce soil nitrate-N content, and how can this be measured? *Plant & Soil* 230, 185-195.
- THORUP-KRISTENSEN,K. 2006. Root growth and nitrogen uptake of carrot, early cabbage, onion and lettuce following a range of green manures. *Soil Use Man* 22, 29-38.
- THORUP-KRISTENSEN,K. & VAN DEN BOOGAARD,R. 1998. Temporal and spatial root development of cauliflower (*Brassica oleracea* L. var. botrytis L.). *Plant & Soil* 201, 37-47.
- THORUP-KRISTENSEN,K. & VAN DEN BOOGAARD,R. 1999. Vertical and horizontal development of the root system of carrots following green manure. *Plant & Soil* 212, 145-153.

7 N uptake

N uptake is calculated as a function of crop N demand on a specific day and the potential root N uptake on the same day. The simulated crop N demand is received from the crop growth part of the model. The potential supply from the soil is calculated as a function of the root length in each soil unit, the content of ammonium-N and nitrate-N in each soil unit and the value S_N read from the croptable parameter file, to control root N uptake efficiency. Diffusion terms are not included in the simulation, as with N uptake this is not assumed to be significant over the relevant time spans for the simulations. N in the form of nitrate is highly mobile in the soil, and diffusion processes will only limit uptake on the very short term even at low root density. Here the equation for potential ammonium-N uptake is shown; this calculation is made for each soil unit and summed for the whole soil volume. The value of S_1 determines the minimum amount of ammonium-N which can be left in the soil:

$$N_{potnh4} = ((\text{rootlength} * S_N * (\text{nnh4} - S_1)) / (S_2 + \text{nnh4})) \quad (4)$$

A minimum level of N left in the soil is included for both ammonium-N and nitrate-N, as experimental data show, that even though crops with high N demand compared to the soil N supply may reduce soil N to very low levels, some soil N is always measured in the soil analyses, especially in the topsoil layers (e.g. Thorup-Kristensen, 2001, 2006). S_2 reduces N uptake as these minimal values are reached.

A function is then used to balance actual N uptake according to crop N demand and potential root N uptake. When crop N demand and potential root N uptake are close to each other, the simulated N uptake will be below either value, but at very high or low

N supply relative to demand, the uptake will be fully controlled by crop N demand and potential root N supply respectively.

$$Nup = Ndemand * (1 - \exp(-1 * (Npot / Ndemand))) \quad (5)$$

Often, the calculated actual N uptake will be lower than the potential root N supply. When this is the case, the actual depletion of soil N will be reduced proportionally from the potential value in all soil units. At last, a specific calculation is made of N taken up from below 0.9 m in the soil. This is made as N leaching loss and other N balance figures are shown mainly from the 0-0.9 m soil layer in much of the model output, and it is therefore necessary also to have an output showing how much N is taken up from below this zone.

Reference List

THORUP-KRISTENSEN, K. 2001. Are differences in root growth of nitrogen catch crops important for their ability to reduce soil nitrate-N content, and how can this be measured? *Plant & Soil* 230, 185-195.

THORUP-KRISTENSEN, K. Effect of deep and shallow root systems on the dynamics of soil inorganic N during three year crop rotations. *Plant & Soil* 288, 233-248. 2006.

Ref Type: Journal (Full)

8 Crop growth and critical N

Each day the increment in plant dry matter is calculated from:

$$\Delta W = \frac{K_2 G_N G_T G_W W}{K_1 + W} \quad (6)$$

where W is the cumulative dry weight, K_2 is calculated as described below. $K_1 = 1 \text{ t ha}^{-1}$. G_T is the effective day degree for the day divided by the average day degree throughout the entire growing period, where the effective day degree is the average temperature for the day less a base temperature, with the limitation that if the average temperature exceeds 20°C then it is set equal to 20°C . G_N and G_W are the growth coefficients dependent on crop %N and water supply. K_2 is calculated from the integral of the above equation with G_N , G_W and G_T set equal to 1. The equation is then

$$K_2 = \frac{K_1 \ln W_{\max} + W_{\max} - K_1 \ln W_P - W_P}{T_h - T_P} \quad (7)$$

where W_P is the dry weight at planting. T_h is the time of final harvest and T_P is the time of drilling or planting in days from Jan 1st.

We use a unified equation to define critical %N for different crops, i.e.

$$\%N_{\text{crit}} = a(1 + be^{-0.26W}) \quad (8)$$

where $\%N_{crit}$ is the critical $\%N$, and a and b are the coefficients, varying from crops (see the crop table of the model).

Luxury N consumption is permitted to take place. It is calculated as follows:

$$\%N_{max} = R_{lux} \%N_{crit} \quad (9)$$

where $\%N_{max}$ is the maximum possible crop $\%N$, and $R_{lux} (\geq 1)$ is the coefficient for luxury N consumption (see the crop table).

For each day a growth coefficient G_N is calculated as:

$$G_N = \min\left(\frac{\%N}{\%N_{crit}}, 1.0\right) \quad (10)$$

where $\%N$ is the actual $\%N$ in the dry matter of the whole plant (excluding fibrous roots)

Similarly, a growth coefficient G_W is calculated as:

$$G_W = \frac{TR_{act}}{TR} \quad (11)$$

where TR and TR_{act} are the actual and potential transpiration rates.

9 Fertility Building Crops

As it is difficult to specify an appropriate target yield for a fertility building crop an alternative approach is used. The user specifies Good, Medium or Bad growth rather than a numerical value and the actual numbers for each crop are read from the crop table. The increment in plant dry matter on each day is calculated from:

$$\Delta W = \min(GG_N G_T W, \Delta W_{type}) \quad (12)$$

where W is the cumulative dry weight, G and ΔW_{type} is set to one of three possible values (good, medium, bad) to define the growth rate and the dry weight increment, G_N and G_T are the growth coefficients dependent on the crop $\%N$ and day degree. The calculation of the growth coefficient G_N is the same as that for a cash crop.

The growth coefficient G_T is calculated:

$$G_T = \begin{cases} 1.0 & \text{if day degree} > 10.0 \\ \frac{\text{day degree} - \text{base temperature}}{10.0 - \text{base temperature}} & \text{base temperature} \leq \text{day degree} \leq 10.0 \\ 0 & \text{day degree} < \text{base temperature} \end{cases}$$

(13)

Another crop parameter specifies the percentage of biomass which is returned to the upper layer of the soil each day; it is then mineralised as a crop residue. This is particularly significant for longer term leys. The user can specify dates at which the crop is mown – on these occasions 50% of the biomass is added to the soil.

Most fertility building crops are legumes and nitrogen fixation is the main source of nitrogen in organic cropping systems. A crop parameter specifies N fixation or not (this also applies to cash crops). The growth of N fixing crops is not limited by nitrogen in the soil as any more that they need is taken from the air.


Annual crops are killed after an appropriate period of time (specified in the Crop Table) regardless of the ‘harvest date’ set by the user. Crops are also killed if the temperature drops below a specified value.

Modelling of the growth of undersown crops begins at the harvest of the overstory with an appropriate dry matter and nitrogen content as specified in the Crop Table; the user can choose between Good, Medium and Bad performance as an understory to give different starting conditions.

10 Economics Module

At present soil- and plant, models rarely contain economic components, because natural and social sciences often use different approaches to modelling. In the EU-Rotate_N model we did not attempt building a separate economic model, rather integrated the economics into a sub-model, so that EU-Rotate_N can run with or without the economic part (Schmutz et. al. 2004). The main entry into the economic model is the total dry matter (TDM), which includes roots, and all above ground dry matter. TDM is an output of the current agronomic model. This parameter however, does not give an indication of the above ground dry matter or fresh matter, nor is there an indication of size or shape of the marketable vegetable parts. Therefore, one of the challenges was finding appropriate algorithms to calculate a marketable yield, which is a major input in any farm economic model. This gives also a figure for the dry matter removed, and the remaining residues (post-harvest) are used as an input for the mineralisation sub-model.

Overview



Simplified model overview and link to economic data

Conversion of total dry matter in marketable yield

Marketable yields are not fixed: the percentage of total yield marketable depends on “soft” or social factors. Among those are market channels, production systems (organic or conventional), eating cultures (e.g. some countries prefer small, other large vegetables, a full-flavoured taste in one may be considered bitter in another). Only a few “hard” figures can be used such as the EU trade classifications, which makes certain vegetables un-marketable if below or above the specifications. Considering these, two strategies were developed - one more empirical the other more theoretical.

(1) For the empirical conversion “Direct conversion” our own research, published and un-published field research data were collected, where both total dry matter and marketable yields were measured across Europe. From this an algorithm was derived converting total dry matter into marketable yield at any given N supply level considering the effects of both sub- and supra-optimal supply of N. A unified algorithm with different crop specific parameters is used for each annual vegetable with a single harvest. There are three main types of vegetable crops: some with a simple constant relationship at all available N levels, some with linear increasing or decreasing relationship depending on available N. Some are more complicated with a non-linear relationship. Other vegetable crops are perennial, like artichokes, or with multiple harvests and need different algorithms than annual, single harvest crops.

“Direct conversion” is a direct conversion of total dry matter (TDM) into marketable yield (MKTY) with one factor R, sourced from empirical data.

$$\text{MKTY} = \text{TDM} \times R \cdot f(N_{av}) \quad (14)$$

With:

- R the ratio of marketable yield to total dry matter for optimum nitrogen supply and spacing
- N_{av} the available nitrogen (N)

The ratio R is individual for each crop and depends on the available N supply used for each crop. The formula for R is a linear or polynomial relationship of available nitrogen (N_{av}).

$$R = r_0 + r_1 \times N_{av} + r_2 \times N_{av}^2 + r_3 \times N_{av}^3 + r_4 \times N_{av}^4 \quad (15)$$

The terms r_0 , r_1 and r_2 are empirically gained for individual crops. For a simple constant relationship: $r_1 = 0$ and $r_2 = 0$. For a linear relationship: $r_2 = 0$. Otherwise, the relationship is non-linear. For some crops, more polynoms may be needed because of different behaviour in sub- and supra-optimum conditions and therefore r_3 , r_4 ... r_x are added.

(2) In a second approach, the single plant fresh weight is calculated. This is done using the harvest index to calculate the dry weight of the harvested parts. Then, with the dry matter content and the plant population, an average single plant fresh weight is produced. Assuming a normal distribution of plant fresh weights and a coefficient of variation of e.g. 20% a lower and upper limit of marketable plant fresh weight can be set (e.g., the EU trade specifications). With this information, an average fresh weight of marketable plants within these specifications is calculated. Using the plant population again, the marketable yield and the residues left post-harvest are calculated.

$$\text{av. single plant fresh weight} = \text{TDM} \times \text{HI} \times \text{plant population}^{-1} \times \%dm^{-1} \quad (16)$$

A normal distribution with a given coefficient of variation is used to simulate the %-gradeout, then the marketable yield in tonnes is calculated. For each crop, a default model choice (approach 1 or 2) is stored in the crop table, but the experienced user can change this.

Prices, variable costs and gross margin calculation

With the marketable yield modelled, the calculation of the crop gross margin (GM) uses the standard equation:

$$GM = \text{MKTY} \times \text{Price} - VC_{ind} - VC_{dep} - VC_{N\ fert} \quad (17)$$

The variable costs independent (VC_{ind}) of marketable yield are recorded per hectare and consist of seed and transplants costs, fertiliser costs excluding N fertiliser, fleece, irrigation, crop protection, weed control. Variable costs depending (VC_{dep}) on the marketable yield are recorded per tonne marketed and multiplied by MKTY. They consist of packaging and drying, transport, harvest casual labour and market commission. The variable costs of inorganic and organic fertilisers ($VC_{N\ fert}$) are calculated using the physical data generated by the model. The triggered amount of fertiliser and number of applications are multiplied by the cost of fertiliser and the cost per application as specified in the input file. Subsidies are not considered in the gross margin calculation. Rotational gross margin is cumulative gross margin of all

crops in the rotation (including the negative gross margin of cover crops) divided by the number of years simulated.

For the calculations, own figures or standardised figures stored in a separate economic data file can be used. In this database, the countries Norway, Denmark, Germany, UK, Italy and Spain are considered. The market channels considered are pre-pack for supermarket, wholesale, direct marketing and processing. The growing systems considered are conventional and organic. The database holds about 300 crop entries of all relevant horticultural crops, including fertility-building crops, across Europe. The data are current prices and standardised variable cost data published in each country for conventional and organic farming systems (e.g. Lampkin et al., 2004; Nix, 2004; Agro Business Consultants Ltd., 2005). The level of data availability and the depth of detail vary among countries.


Literature cited.

Agro Business Consultants Ltd. (2005): The Agricultural Budgeting & Costing Book, Agro Business Consultants.


Lampkin, N., M. Measures, et al. (2004). 2004 Organic Farm Management Handbook. Aberystwyth, Newbury, Organic Farming Research Unit at Institute of Rural Sciences, University of Wales. Organic Advisory Service, Elm Farm Research Centre. 226.

Nix, J. (2004). Farm Management Pocketbook 2005. London, The Andersons Centre.

Schmutz, U., C. Firth, F. Rayns and C. Rahn (2004): Can N Use and Farm Income be Optimized for Organic Field Vegetable Rotations in Europe? In: Organic Farming Ed. A. Hopkins BGS Symposium No. 37: ISBN 0905944 844. p. 200-203



FIGUUR 1: EVOLUTIE VAN HET KOOLSTOFGEHALTE VAN DE BODEM VOOR DE BRAAKBEHANDELINGEN (0 – 60 – 120 KG N HA⁻¹).



FIGUUR 2: EVOLUTIE VAN HET KOOLSTOFGEHALTE VAN DE BODEM VOOR GELE MOSTERD BIJ VROEGE ZAAI (0 – 60 – 120 KG N HA⁻¹).